

The year in cardiovascular medicine 2021: heart failure and cardiomyopathies

**Johann Bauersachs^{1*}, Rudolf A. de Boer², Joann Lindenfeld³,
and Biykem Bozkurt⁴**

¹ Department of Cardiology and Angiology, Hannover Medical School, Carl-Neuberg-Straße 1, 30625 Hannover, Germany

² Department of Cardiology, University Medical Center Groningen, Groningen, The Netherlands

³ Vanderbilt Heart and Vascular Institute, Vanderbilt University Medical Center, Nashville, TN, USA

⁴ Winters Center for Heart Failure, Cardiology, Baylor College of Medicine and Michael E. DeBakey VA Medical Center, Houston TX, USA

Received Date: 16 September 2021; Revision received Date: 27 October 2021; Accepted Date: 16 November 2021

Reproduced from: Bauersachs J, de Boer RA, Lindenfeld J, Bozkurt B. The year in cardiovascular medicine 2021: heart failure and cardiomyopathies. *Eur Heart J* 2022;43:367–376. <https://doi.org/10.1093/eurheartj/ehab887>, by permission of Oxford University Press on behalf of the European Society of Cardiology

© The Author(s) 2022.


All rights reserved; no part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without prior written permission of the Publishers.

For Permissions, please email: journals.permissions@oup.com

The opinions expressed in the Journal item reproduced as this reprint are those of the authors and contributors, and do not necessarily reflect those of the European Society of Cardiology, the editors, the editorial board, Oxford University Press or the organization to which the authors are affiliated.

The mention of trade names, commercial products or organizations, and the inclusion of advertisements in this reprint do not imply endorsement by the Journal, the editors, the editorial board, Oxford University Press or the organization to which the authors are affiliated. The editors and publishers have taken all reasonable precautions to verify drug names and doses, the results of experimental work and clinical findings published in the Journal. The ultimate responsibility for the use and dosage of drugs mentioned in this reprint and in interpretation of published material lies with the medical practitioner, and the editors and publisher cannot accept liability for damages arising from any error or omissions in the Journal or in this reprint. Please inform the editors of any errors.

Oxford University Press, OPL, and the European Society of Cardiology are not responsible or in any way liable for the accuracy of the reprint, for any errors, omissions, or inaccuracies, or for any consequences arising therefrom. The Czech Society of Cardiology is solely responsible for this reprint.



Graphical Abstract Summary of the universal definition and EF classification of heart failure; management of HFrEF according to 2021 ESC guidelines for heart failure and results of the EMPEROR-preserved trial.

Abstract

In the year 2021, the universal definition and classification of heart failure (HF) was published that defines HF as a clinical syndrome with symptoms and/or signs caused by a cardiac abnormality and corroborated by elevated natriuretic peptide levels or objective evidence of cardiogenic congestion. This definition and the classification of HF with reduced ejection fraction (HFrEF), mildly reduced, and HF with preserved ejection fraction (HFpEF) is consistent with the 2021 ESC Guidelines on HF. Among several other new recommendations, these guidelines give a Class I indication for the use of the sodium–glucose co-transporter 2 (SGLT2) inhibitors dapagliflozin and empagliflozin in HFrEF patients. As the first evidence-based treatment for HFpEF, in the EMPEROR-Preserved trial, empagliflozin reduced the composite endpoint of cardiovascular death and HF hospitalizations. Several reports in 2021 have provided novel and detailed analyses of device and medical therapy in HF, especially regarding sacubitril/valsartan, SGLT2 inhibitors, mineralocorticoid receptor antagonists, ferric carboxymaltose, soluble guanylate cyclase activators, and cardiac myosin activators. In patients hospitalized with COVID-19, acute HF and myocardial injury is quite frequent, whereas myocarditis and long-term damage to the heart are rather uncommon.

Keywords: Heart failure; Epidemiology; Imaging; Biomarkers; Pharmacotherapy; Artificial intelligence; SGLT-2 inhibitor; Angiotensin receptor–neprilysin inhibitors; Activators of soluble guanylate cyclase; Device therapy

Introduction

Heart failure (HF) remains a major challenge for patients and healthcare systems worldwide. For patients suffering from HF with reduced ejection fraction (HFrEF), several evidence-based treatments are available and have markedly improved prognosis and quality of life; however, a subset of these patients displays a rapid progression of HF despite best care. A recent special article called to action for global approaches to novel drug solutions for these patients,¹ but also for patients with HF with preserved EF (HFpEF), for whom until recently there was not a single evidence-based treatment.

In this article, we summarize important progress that has been made in 2021 regarding the diagnosis and treatment of HF with a special focus on articles published in 2021 in the *European Heart Journal* and the *European Journal of Heart Failure*.

Definition and classification of heart failure

With the recognition of the need for standardization of an HF definition, the Universal Definition and Classification of Heart Failure was developed, which defined HF as a clinical syndrome with current or prior symptoms and/or signs caused by a structural and/or functional cardiac abnormality and corroborated by elevated natriuretic peptide (NP) levels or objective evidence of cardiogenic pulmonary or systemic congestion by diagnostic modalities (Figure 1).² It also provided revised definitions for stages of HF, categorized as 'At-Risk for HF' (former Stage A) for patients at risk for HF but without current or prior symptoms or signs of HF and without structural cardiac changes or elevated biomarkers of heart disease; Pre-HF (former Stage B) for patients without current or prior symptoms or signs of HF but evidence of structural heart disease, abnormal cardiac function, elevated NP levels or elevated cardiac troponin levels; 'Heart Failure' (former Stage C) for symptomatic patients, 'Advanced HF' (former Stage D) for patients with severe symptoms and/or signs of HF (Figure 1 – see in original). Ejection fraction categories were classified as HFrEF: left ventricular (LV) EF $\leq 40\%$ (Figure 1 – see in original); HF with mildly reduced EF (HFmrEF): LVEF 41–49%; HFpEF: LVEF $>50\%$; and HF with improved EF (HFimpEF): HF with a baseline LVEF $\leq 40\%$, a ≥ 10 point increase from baseline LVEF, and a second measurement of LVEF $>40\%$. The EF categories used in the recent 2021 ESC HF Guidelines were consistent with these classifications.³ In the Universal Definition of HF, there was also an emphasis on trajectories of HF and to use 'persistent HF' instead of 'stable HF' for patients with ongoing symptoms/signs and 'HF in remission' instead of 'recovered HF' for patients with resolution of symptoms and signs of HF or with the resolution of previous structural/functional heart disease² (Figure 1 – see in original). Though a simple definition of HF predominantly depending on NPs was proposed as an alternative,⁴ limitations of such an approach due to variability of NP levels by age, sex, body mass, renal function, and atrial fibrillation; and lack of

specificity and lack of evidence in linking treatments to a biomarker-based approach were identified as significant barriers to a simply biomarker-based approach in definition of HF.⁴

Epidemiology

The HF Atlas survey reports a wide-ranging incidence of HF and HF hospitalizations across Europe with considerable heterogeneity in the resources for management and the data quality providing data to allow the development of strategies to improve inequalities.⁵ Exposure to ambient air pollutants increases the risk of HF in a dose-dependent fashion, and there was a particularly high risk of HF among persons with genetic higher susceptibility to HF (Figure 2).⁶ Air pollution probably should be considered in risk scores to predict HF.

A recent European registry report demonstrated that dilated cardiomyopathy (DCM), not skeletal myopathy, is the major determinant of prognosis in patients with dystrophin gene mutations.⁷ Finally, cancer and HF occur more commonly together than predicted by risk models, and a recent study suggests that statins reduce the risk of both and have a greater risk reduction with more prolonged use.⁸

Diagnostics and risk stratification

For HFrEF, the main diagnostic criterion remains LVEF $\leq 40\%$.³ However, there is more controversy in the other categories, HFmrEF and HFpEF. Pieske et al.⁹ formulated, on behalf of the ESC, new diagnostic criteria, including echo parameters, NPs, and if a definitive diagnosis cannot be made, to turn to stress testing and/or invasive haemodynamics.

There is increasing appreciation that classical diagnostics fall short in complex multifactorial diseases with various aetiologies and precipitants, and several studies have addressed whether an agnostic approach, where large data sets are queried by computer algorithms, may be superior in making a specific diagnosis. Such techniques are referred to as machine learning (ML) and artificial intelligence (AI). Peyster et al.¹⁰ used an automated image analysis to detect rejection after heart transplantation and described a 'Computer-Assisted Cardiac Histologic Evaluation (CACHE)-Grader' pipeline that was non-inferior to the rejection grading provided by independent pathologists. Another field of research for which AI provides an attractive tool is the categorization of patients who received a general diagnosis of HF. Verdonschot et al.¹¹ studied 795 consecutive DCM patients with data on aetiology and co-morbidities, imaging studies and endomyocardial biopsies, and identified four distinct phenogroups. Woolley et al.¹² using an algorithm based on 363 biomarkers to phenotype, 429 patients with HFpEF identified four clusters with different clinical parameters and important differences in prognosis.

Artificial intelligence/machine learning might be particularly useful for a diagnosis of HF. Kwon et al.¹³ evalu-




Figure 2 Long-term joint exposure to various air pollutants, including $PM_{2.5}$, PM_{10} , $PM_{2.5-10}$, NO_2 , and NO_x is associated with an elevated risk of incident heart failure in an additive manner. Persons with genetic higher susceptibility to heart failure displayed a particularly high risk of heart failure. Reprinted with permission from Wang *et al.*,⁶ by permission of OUP on behalf of ESC.

ated data from 34 103 patients who underwent echocardiography and electrocardiogram (ECG) and created an ML algorithm that could detect HFpEF. Segar *et al.*¹⁴ employed ML models to aid in predicting race-specific risk for incident HF.

In the near future, we will be faced with many more potential utility of AI/ML models, as there is a clear need for individualized approaches and decision-making.¹⁵ It will be essential, however, to provide recommendations as to what input is (minimally) required for models, and the models must be prospectively tested in independent settings. Furthermore, treatment decisions based on the models must be tested in a randomized blinded fashion.¹⁶

Imaging and biomarkers

A state-of-the-art diagnosis of HF remains challenging. The ESC guidelines³ recommend using an array of signs and symptoms, supplemented with imaging and biomarkers studies. The imaging primarily relies on echocardiography and CMR, and NPs and high sensitivity troponins are the preferred biomarkers. However, sophisticated classification of patients in various categories using imaging and biomarkers may enhance adequate phenotyping,^{11,17} and imaging of non-cardiac tissues such as fat may have relevance to HF phenotyping, too.^{18,19} Furthermore, next-generation genetic analyses has been shown to have a consequence for prognosis²⁰ and diagnosis²¹ of HF. In addition, a recent

article highlighted the indications of endomyocardial biopsies.²²

Specific situations

Acute heart failure

The 2021 ESC guidelines did not significantly change recommendations for acute HF, although the use of opioids was downgraded to a Class III recommendation.³ Evidence continues to accrue supporting the use of urinary sodium in assessing outcomes in acute HF.^{23,24}

Cardiogenic shock

Mortality remains high in cardiogenic shock, and randomized trials assessing therapies remain rare but a single-centre trial randomized patients with cardiogenic shock to either milrinone or dobutamine and showed no differences in any of the primary or secondary outcomes.²⁵ In the follow-up of the IMPRESS trial in cardiogenic shock, there was no difference in mortality comparing intra-aortic balloon pumps vs. the Impella device at 5 years.²⁶ A biomarker composite outperformed other risk scores for cardiogenic shock using 4 biomarkers [Cystatin C, Lactate, interleukin-6, and N-terminal pro brain natriuretic peptide (NT-proBNP)].²⁷ A recent consensus statement outlines important suggestions for optimizing cardiogenic shock trials.²⁸

Ventricular assist devices and heart transplantation

A single entry registry confirms that HeartMate III (HMIII) outcomes are better than historical controls confirming randomized trials.²⁹ The stroke rate with HMIII is less than with the Heartware ventricular assist device (HVAD)—one of several reasons the HVAD has been withdrawn from use.³⁰ Disappointingly, left ventricular assist devices (LVAD) use does not reduce myocardial fibrosis nor does a new risk score improve the prediction of right ventricular failure post-LVAD, but on the bright side, elderly patients have benefits in quality of life and exercise capacity with LVADs.^{31–33} There is substantial inter-observer variability in the diagnosis of cellular rejection in myocardial biopsies but automated computation image analysis may allow improved standardization as described in the section on Diagnostics and Imaging. Non-invasive prediction of rejection in cardiac transplant recipients has been elusive, but studies using peripheral blood cell-free DNA show promising early results.³⁴

Pregnancy/patients with peripartum cardiomyopathy

Women with a known cardiomyopathy or at risk for HF planning pregnancy, or presenting with HF during or after pregnancy are in need of individualized pre-, during, and post-pregnancy assessment and counselling.³⁵

Patients with peripartum cardiomyopathy are at risk for detrimental outcomes^{36,37} but often do recover from HFrEF. Recent publications investigated the value of ECG abnormalities for predicting echocardiographic results and the role of hypertensive disorders during pregnancy.^{38,39}

Hypertrophic cardiomyopathy/amyloidosis

In the health status analysis of EXPLORER-HCM, mavacamten markedly improved the health status of patients with symptomatic obstructive hypertrophic cardiomyopathy (HCM) compared with placebo.⁴⁰ Gaps in evidence for risk stratification for sudden cardiac death in HCM were summarized by Pelliccia et al.⁴¹ In a study by Marston et al.⁴² using Sarcomeric Human Cardiomyopathy Registry, patients with childhood-onset HCM were reported more likely to have sarcomeric disease, carry a higher risk of life-threatening ventricular arrhythmias, and have a greater need for advanced HF therapies. In the German Cardiac Society position statement, Yilmaz et al.⁴³ outline a diagnostic algorithm to detect cardiac amyloidosis, to accurately determine its extent, and to reliably identify the underlying subtype of amyloidosis, thereby enabling subsequent targeted treatment.

Cancer

Heart failure often complicates the treatment of cancer, and a recent paper proposes definitions of cardiovascular (CV) toxicities.⁴⁴ Classically, chemotherapy and radiotherapy have been identified as risk factors, but in the recent decade, immunotherapy with immune checkpoint inhibitors (ICIs) is becoming the mainstay of cancer treatment. However, ICIs also carry a risk for CV side effects. D'Souza et al.⁴⁵ reported on this risk in a Danish registry and show that ICI is associated with a 1.8% 1-year risk for (peri-) myocarditis, and with an almost 10% risk for any CV complication. Given the increasing use of ICI, this issue will require clinical guidance and further study, as ICIs have an impact on several cells and tissues.^{46,47} There are initial reports providing guidance as to treat ICI-induced myocarditis.^{48,49}

This field extends the increasing awareness that incident cancer is more common in patients with prevalent HF,⁵⁰ and that cancer and HF may be connected more closely than anticipated before. In support of this, Ren et al.⁸ demonstrated that the use of statins reduces incident cancer. Finally, a special article by Zannad et al.⁵¹ discusses aspects of cancer research that may be applicable to HF research, with the aim of streamlining the clinical trial process and decreasing the time and cost required to bring safe, effective, treatments to HF patients.

Pharmacotherapies

New algorithm of the 2021 ESC Guidelines on heart failure for the pharmacological treatment of heart failure with reduced ejection fraction

The 2021 ESC Guidelines on HF provide a Class I recommendation for pharmacological treatment of all HFrEF patients with a combination of an angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor-neprilysin inhibitor (ARNI), a beta-blocker, a mineralocorticoid receptor antagonist (MRA), and a sodium–glucose co-transporter 2 (SGLT2) inhibitor (dapagliflozin or empagliflozin) (Figure 1B – see in original).³ The guideline still recommends the use of ARNI as a replacement for ACE inhibitor; however, an ARNI may also be considered as a first-line therapy instead of an ACE inhibitor.

It is recommended that these four disease-modifying drugs are initiated within a short time frame.^{3,52} Potential advantages of another algorithm for the sequencing of these drugs have been suggested by McMurray and Packer⁵³ with beta-blockade and SGLT2 inhibition as first-line therapies. However, albeit appealing from a pathophysiological standpoint such a new sequence is not yet evidence-based.

A recent consensus document of the HFA of the ESC identified nine patient profiles that may be relevant for treatment implementation in patients with HFrEF taking into account heart rate, atrial fibrillation, symptomatic low blood pressure, estimated glomerular filtration rate, or hyperkalaemia. Using such a personalized approach may lead to a better and more comprehensive therapy for each individual patient.⁵⁴

Angiotensin-converting enzyme inhibition

While ACE inhibitors are a standard for the prevention and treatment of HF for many years, the impact of these drugs as preventive therapy for HF in patients with Duchenne muscular dystrophy was unclear. A large French registry showed that prophylactic treatment of patients without LV dysfunction with an ACE inhibitor was able to prevent the transition to HF and improve survival in Duchenne muscular dystrophy.⁵⁵

Angiotensin receptor-neprilysin inhibitors (PARAGON, PARADIGM, PARALLAX, PARADISE-MI, LIFE)

In an analysis of the PARADIGM-HF trial, initiation of sacubitril/valsartan, even when titrated to target dose, did not lead to greater discontinuation or down-titration of other guideline-directed medical therapies and was associated with fewer discontinuations of MRA.⁵⁶ In real-world patients with HFrEF, sacubitril/valsartan was effective, safe, and well tolerated.^{57–60} Sacubitril-valsartan was found to be useful in treating resistant hypertension in HFpEF in the PARAGON-HF trial when compared with valsartan.⁶¹ In the PROVE-HF trial, in patients with HFrEF, 32% improved their EF to >35% by 6 months and 62% to >35% by 12 months after initiation of sacubitril/valsartan therapy.⁶² In patients with asymptomatic LV systolic dysfunction late after myocardial infarction, treatment with sacubitril/valsartan did not have a significant reverse remodelling effect compared with valsartan.^{63,64} In the PARADISE-MI trial,⁶⁵ sacubitril/valsartan did not significantly reduce the rate of CV death, HF hospitalization, or outpatient HF requiring treatment in patients with LVEF ≤40% and/or pulmonary congestion following acute myocardial infarction, compared with ramipril (results presented at the ACC). In the Sacubitril/Valsartan in Patients with Advanced Heart Failure with Reduced Ejection Fraction in the Advanced Heart Failure (LIFE-HF) trial, which enrolled NYHA Class IV patients and LVEF ≤35%, sacubitril/valsartan did not improve the clinical composite endpoints (presented at ACC 2021). PARALLAX trial will determine if sacubitril/valsartan improves NT-proBNP levels, exercise capacity, quality of life, and symptom burden in HF patients with EF >40%.⁶⁶

In the new 2021 ESC Guidelines on HF,³ sacubitril/valsartan is recommended as a replacement for an ACE inhibitor in patients with HFrEF as a Class I recommen-

dation. Initiation of sacubitril/valsartan in ACE inhibitor naive patients with HFrEF on the other hand is suggested as a Class IIb recommendation.³

Sodium-glucose co-transporter 2 inhibitors (EMPEROR-Reduced, EMPEROR-Preserved, DAPA-HF, SOLOIST)

Sodium-glucose co-transporter 2 inhibitors are rapidly becoming the panacea for the entire spectrum of cardio-metabolic and renal disease. In trials in type 2 diabetes mellitus (T2DM), a beneficial effect was observed for CV endpoints in general, while the effects on incident HF were overwhelmingly positive. These effects were validated in patients with prevalent HFrEF, first in DAPA-HF and a year later in the EMPEROR-Reduced trial. Numerous subanalyses from these trials were published in 2021.

First, besides the striking effects on hard endpoints, it is more and more recognized that functional status and symptoms are important to patients with HFrEF.⁶⁷ Both in DAPA-HF and EMPEROR-Reduced, these were improved,^{68,69} although a smaller dedicated trial with empagliflozin did not improve functional status.⁷⁰ Further, a series of subanalyses showed no interaction of SGLT2 inhibitors with common HF drugs, such as MRAs, and most importantly, also not with sacubitril/valsartan.^{71,72} Furthermore, the equal effects of the drugs were ascertained by analysing the effects across countries and ethnicities.⁷³ Another striking observation was that dapagliflozin was associated with a lower incidence of new-onset diabetes.⁷⁴ Collectively, to date, we have not seen any analysis suggesting a differential or lesser effect of SGLT2 inhibitors in HFrEF. We therefore must start to learn how to employ these drugs practically.^{52,75}

Different from HFrEF, the efficacy of SGLT2 inhibitors in HFpEF remained to be proven. However, the EMPEROR-Preserved study presented during ESC 2021 demonstrated that empagliflozin reduced the primary combined endpoint of CV death and HF hospitalization in almost 6000 patients with HFpEF (*Figure 3 – see in original*). These data are extremely important and provide hope for millions of HFpEF patients for whom there were no evidence-based therapies. Over a median follow-up of 26 months, the primary outcome event occurred in 13.8% of the patients in the empagliflozin group and in 17.1% in the placebo group [hazard ratio (HR): 0.79; 95% confidence interval (CI): 0.69–0.90; $P < 0.001$]. Empagliflozin was very effective in reducing HF hospitalization, but all-cause mortality was not reduced. The effects of empagliflozin were consistent in patients with or without diabetes.^{76,77} Shortly, the result of the second mortality trial in HFpEF with the SGLT2 inhibitor dapagliflozin, DELIVER, will be presented.⁷⁸

Sodium-glucose co-transporter 2 inhibitors were also evaluated in patients with acute HF or immediately after acutely decompensated HF. The SOLOIST trial,⁷⁹ with the mixed SGLT 1/2 inhibitor sotagliflozin, enrolled 1244 patients with T2DM and recent worsening HF and showed a beneficial effect of the study drug, initiated before or shortly after discharge, with regard to a significantly lower total number of CV deaths and HF hospitalizations and urgent visits for HF. The ongoing EMPULSE trial will provide more data in the acute HF arena.⁸⁰

Sodium–glucose co-transporter 2 inhibitors do not stop to amaze us in renal disease. After the publication of the hallmark trials CREDENCE and DAPA-CKD,⁸¹ in 2021, the SCORED trial⁸² came out, demonstrating in patients with T2DM and chronic kidney disease, allocated to sotagliflozin or placebo, a reduction of 37% in the primary endpoint of CV death and HF events (HR: 0.74; 95% CI: 0.63–0.88; $P < 0.001$). However, sotagliflozin was associated with adverse events such as diarrhoea, genital mycotic infections, volume depletion, and diabetic ketoacidosis.

Mineralocorticoid receptor antagonists (FIDELIO, FIGARO, HOMAGE)

Mineralocorticoid receptor antagonists are first-line therapies for HFrEF and may also be considered in HFmrEF.³ Novel non-steroidal MRA such as finerenone differ from steroidal MRA regarding tissue distribution, MR binding, recruitment of cofactors, and downstream gene expression.⁸³ In FIDELIO-DKD, finerenone improved CV and kidney outcomes in patients with chronic kidney disease and T2D regardless of baseline HF status (G. Filippatos, 2021, submitted for publication). In FIGARO-DKD, finerenone reduced the primary composite endpoint of death from CV causes, non-fatal myocardial infarction, non-fatal stroke, or HF hospitalization with the benefit driven primarily by a lower incidence of HF hospitalization.⁸⁴ In HOMAGE, in patients with, or at high risk for, coronary disease and raised NP levels, no interaction between baseline serum galectin-3 and changes in procollagen collagen biomarkers induced by spironolactone treatment was observed. However, blood pressure and NT-proBNP were reduced by spironolactone.⁸⁵

Activators of soluble guanylate cyclase (VICTORIA)

The novel activator of soluble guanylate cyclase, vericiguat, in a subanalysis of the VICTORIA trial, did not reduce new-onset atrial fibrillation. However, pre-existing atrial fibrillation did not affect the beneficial effect of vericiguat on the primary composite outcome (time to CV death or first HF hospitalization) or its components.⁸⁶ Similarly, beneficial effects of vericiguat were consistent across the full range of renal function.⁸⁷

Cardiac myosin activators

A substudy of the pivotal trial of the myosin activator omecamtiv mecarbil (GALACTIC-HF) in patients with HFrEF found that the drug reduced the primary endpoint of HF hospitalization and CV death more as EF declined with a 17% decrease in the lowest quartile ($EF \leq 22\%$) and no benefit in the highest quartile ($EF \geq 33\%$).⁸⁸

Ferric carboxymaltose (AFFIRM-AHF; IRON-CRT)

Iron deficiency is related to worse outcomes in HF. The AFFIRM-AHF study demonstrated that in patients with LVEF $< 50\%$ and iron deficiency after a hospitalization for acute HF, i.v. treatment with ferric carboxymaltose did not only reduce HF hospitalizations but also results in clinically meaningful beneficial effects on quality of life.⁸⁹ In HFrEF patients with iron deficiency and a persistently

reduced LVEF $< 45\%$ after cardiac resynchronization therapy (IRON-CRT) study, i.v. ferric carboxymaltose FCM improved cardiac structure and function, as well as quality of life.⁹⁰

Iron deficiency also contributes to resistance to endogenous erythropoietin, an important cause of anaemia in HF.⁹¹

Others

In a small clinical trial, CDR132L, an antisense oligonucleotide drug directed against miR-132 was well tolerated and seemed to be associated with cardiac functional improvement in HF patients.^{92–94}

In 50 patients with idiopathic chronic DCM and parvovirus B19 persistence, i.v. immunoglobulin therapy did not significantly improve LV systolic function or functional capacity beyond standard medical therapy.⁹⁵

Device and interventional therapies

Cardiac resynchronization therapy

In patients with HF, atrial fibrillation and a narrow QRS mortality and HF hospitalizations were reduced by atrioventricular junctional ablation and cardiac resynchronization therapy (CRT) compared with pharmacological treatment alone; this beneficial effect was similar in patients with LVEF $\leq 35\%$ and $> 35\%$.⁹⁶ Guidelines for CRT and suggestions for optimized implementation have recently been published.^{97,98} The controversy about whether adding an ICD to CRT provide additional mortality benefit, especially in non-ischaemic HF continues.⁹⁹

Percutaneous mitral valve repair

The US Valvular Disease Guidelines as well as the 2021 ESC Guidelines on valvular heart disease recently upgraded the recommendation for transcatheter mitral valve repair (TEER) for secondary (functional) mitral regurgitation (SMR) to a IIA recommendation for patients who meet COAPT criteria.^{100,101} A joint position statement from the ESC supports this recommendation.¹⁰² The 3-year results of the COAPT trial demonstrate the ongoing benefit of TEER.¹⁰³ An important secondary analysis from COAPT demonstrates that residual 3–4+ SMR is the strongest risk factor for poor outcomes in both the TEER group and in the medical therapy group.¹⁰⁴ In patients with atrial fibrillation, TEER was associated with a lower risk of stroke.¹⁰⁵ Subgroups of MITRA-FR mimicking COAPT patients did not show a benefit of TEER, although a subgroup of COAPT mimicking MITRA-FR patients did show a benefit in HF hospitalizations.^{106,107}

Implantable haemodynamic monitors

The GUIDE-HF trial evaluated haemodynamic guided management to reduce HF hospitalizations and mortality in patients with NYHA II–IV and all ejection fractions. The overall analysis was negative but when COVID-19 was accounted for there was a significant reduction in HF hospitalization in NYHA II–III patients with either a previous HF hospitalization or elevated NPs.¹⁰⁸

Specific management

Telemedicine and remote monitoring

In a comprehensive review, Bekfani and colleagues discuss unmet needs in the management of patients with HF, how remote monitoring might contribute to future solutions and provide an overview of current and novel remote monitoring technologies.¹⁰⁹ A great variety of innovative remote monitoring technologies and algorithms including patient self-managed testing, wearable devices, technologies integrated into clinically indicated therapeutic devices, such as pacemakers and defibrillators, and landmark clinical trials of remote monitoring were reviewed.

Rehabilitation

In an Expert Panel consensus document on Cardiac Rehabilitation for Patients with Heart Failure, Bozkurt *et al.*¹¹⁰ provide an overview of efficacy and safety evidence of exercise training and cardiac rehabilitation in HFrEF and HFpEF, recommendations on practical approaches to exercise training and cardiac rehabilitation in patients with HF

and examine the reasons and solutions for underutilization of cardiac rehabilitation in HF patients. In the REHAB-HF trial, in a diverse population of older patients who were hospitalized for acute decompensated HF, an early, transitional, tailored, progressive rehabilitation intervention that included multiple physical function domains resulted in greater improvement in physical function than usual care. This is an important study demonstrating the safety and efficacy of initiation of progressive rehabilitation initiated during and early post-hospitalization in HF patients regardless of LVEF.¹¹¹

Heart failure during the COVID-19 pandemic

Incident acute HF was recognized as a complication in 2%, and myocardial injury in 10% of all patients hospitalized with COVID-19.¹¹² Elevated admission NT-proBNP levels were associated with higher mortality,¹¹³ and cardiac myocyte-specific microRNAs were upregulated in critically ill COVID-19 patients indicating cardiac involvement.¹¹⁴




Figure 4 Myocardial injury in recovered COVID-19 patients assessed by cardiovascular magnetic resonance. Myocarditis-like injury can be encountered, with limited extent and minimal functional consequence. Reprinted with permission from Kotecha *et al.*,¹²⁷ by permission of OUP on behalf of ESC.

Declining overall admission rates for HF¹¹⁵ and higher out-of-hospital mortality rates¹¹⁶ during lockdown were recognized as alarming issues, reflecting lack of access to care among patients with established HF. Randomized trials demonstrated the safety of continuation of ACE inhibitors or ARB among patients hospitalized with COVID-19.^{117–119} Dapagliflozin treatment did not significantly reduce organ dysfunction or death, but was well tolerated in patients hospitalized with COVID-19 (DARE-19 trial).¹²⁰ Myocarditis emerged as a rare complication of COVID-19 mRNA vaccinations, especially in young men.¹²¹ Benefit-risk assessment for COVID-19 vaccination was favourable for all age and sex groups; and almost all patients with myocarditis had resolution of symptoms and signs.¹²¹ Long-term complications of SARS-CoV-2 infection include persistent sinus tachycardia, postural orthostatic tachycardia syndrome, atrial arrhythmia, and cardiomyopathy.¹²² Among athletes recovering from COVID-19, several CMR studies reported varying rates and degrees of cardiac abnormalities suggestive of myocarditis.^{123,124} Screening by troponin, ECG, echocardiography, and additional CMR and/or stress echocardiography if abnormal, resulted in only 0.6% of the athletes being restricted to return to sports, and none had cardiac events.¹²⁵ Though myocardial injury is common in COVID-19, and SARS-CoV-2 RNA can be detected in the heart, myocarditis is an uncommon pathologic diagnosis occurring in 4.5% of highly selected cases undergoing autopsy or endomyocardial biopsy.¹²⁶ During convalescence after severe COVID-19 infection with troponin elevation, myocarditis-like injury can be detected by CMR, however, with limited extent and minimal functional consequence (Figure 4).¹²⁷

Acknowledgements

The authors wish to thank Dr T. König for editorial assistance.

Conflict of interest: J.B. received honoraria for lectures/consulting from Novartis, Vifor, Bayer, Servier, Abiomed, Pfizer, Boehringer Ingelheim, AstraZeneca, Cardior, Daiichi Sankyo, CVRx, BMS, MSD, Amgen, Corvia, not related to this article; and research support for the department from Zoll, CVRx, Vifor, Abiomed, not related to this article. In addition, J.B. is Scientific Advisory Board Member of Cardior and has a patent PCT/EP2007/008772 with royalties paid, and a patent PCT/EP2009/051986 with royalties paid both on microRNA and downstream targets for diagnostic and therapeutic purposes. The UMCG, which employs R.A.d.B. has received research grants and/or fees from AstraZeneca, Abbott, Boehringer Ingelheim, Cardior Pharmaceuticals GmbH, Ionis Pharmaceuticals, Inc., Novo Nordisk, and Roche. R.A.d.B. received speaker fees from Abbott, AstraZeneca, Bayer, Novartis, and Roche. J.L. received honoraria for consultation from Abbott, Alleviant, AstraZeneca, Boehringer Ingelheim, Boston Scientific, CVRx, Edwards Lifesciences, Merck, Vwave and grant function from AstraZeneca, Volumetrix, Sensible Medical. B.B. received honoraria for consultation/advisory committee participation from Bayer, scPharmaceuticals, Vifor Pharma, Amgen, Relpysa, Baxter; and served in the Clinical Events Committee for Guide-HF Trial Abbott Pharmaceuticals and Data Safety Monitoring Board for Anthem Trial by Liva Nova Pharmaceuticals; none related to this article.

References

1. Figtree GA, Broadfoot K, Casadei B, Califf R, Crea F, Drummond GR, et al. A call to action for new global approaches to cardiovascular disease drug solutions. *Eur Heart J* 2021;42:1464–1475.
2. Bozkurt B, Coats AJS, Tsutsui H, Abdelhamid CM, Adamopoulos S, Albert N, et al. Universal definition and classification of heart failure: a report of the Heart Failure Society of America, Heart Failure Association of the European Society of Cardiology, Japanese Heart Failure Society and Writing Committee of the Universal Definition of Heart Failure: endorsed by the Canadian Heart Failure Society, Heart Failure Association of India, Cardiac Society of Australia and New Zealand, and Chinese Heart Failure Association. *Eur J Heart Fail* 2021;23:352–380.
3. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Bohm M, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J* 2021;42:3599–3726.
4. Cleland JGF, Pfeffer MA, Clark AL, Januzzi JL, McMurray JJV, Mueller C, et al. The struggle towards a Universal Definition of Heart Failure—how to proceed? *Eur Heart J* 2021;42:2331–2343.
5. Seferovic PM, Vardas P, Jankowska EA, Maggioni AP, Timmis A, Milinkovic I, et al. The heart failure association atlas: heart failure epidemiology and management statistics 2019. *Eur J Heart Fail* 2021;23:906–914.
6. Wang M, Zhou T, Song Y, Li X, Ma H, Hu Y, et al. Joint exposure to various ambient air pollutants and incident heart failure: a prospective analysis in UK Biobank. *Eur Heart J* 2021;42:1582–1591.
7. Restrepo-Cordoba MA, Wahbi K, Florian AR, Jimenez-Jaimez J, Politano L, Arad M, et al. Prevalence and clinical outcomes of dystrophin-associated dilated cardiomyopathy without severe skeletal myopathy. *Eur J Heart Fail* 2021;23:1276–1286.
8. Ren Q-W, Yu S-Y, Teng T-HK, Li X, Cheung K-S, Wu M-Z, et al. Statin associated lower cancer risk and related mortality in patients with heart failure. *Eur Heart J* 2021;42:3049–3059.
9. Pieske B, Tschöpe C, de Boer RA, Fraser AG, Anker SD, Donal E, et al. How to diagnose heart failure with preserved ejection fraction: the HFA-PEFF diagnostic algorithm: a consensus recommendation from the Heart Failure Association (HFA) of the European Society of Cardiology (ESC). *Eur Heart J* 2019;40:3297–3317.
10. Peyster EG, Arabyarmohammadi S, Janowczyk A, Azarianpour-Esfahani S, Sekulic M, Cassol C, et al. An automated computational image analysis pipeline for histological grading of cardiac allograft rejection. *Eur Heart J* 2021;42:2356–2369.
11. Verdonschot JAJ, Merlo M, Dominguez F, Wang P, Henkens M, Adriaens ME, et al. Phenotypic clustering of dilated cardiomyopathy patients highlights important pathophysiological differences. *Eur Heart J* 2021;42:162–174.
12. Woolley RJ, Ceelen D, Ouwerkerk W, Tromp J, Figarska SM, Anker SD, et al. Machine learning based on biomarker profiles identifies distinct subgroups of heart failure with preserved ejection fraction. *Eur J Heart Fail* 2021;23:983–991.
13. Kwon J-M, Kim K-H, Eisen HJ, Cho Y, Jeon K-H, Lee SY, et al. Artificial intelligence assessment for early detection of heart failure with preserved ejection fraction based on electrocardiographic features. *Eur Heart J Digital Health* 2021;2:106–116.
14. Segar MW, Jaeger BC, Patel KV, Nambi V, Ndumele CE, Correa A, et al. Development and validation of machine learning-based race-specific models to predict 10-year risk of heart failure: a multicohort analysis. *Circulation* 2021;143:2370–2383.
15. Hamdani N, Costantino S, Mugge A, Lebeche D, Tschöpe C, Thum T, et al. Leveraging clinical epigenetics in heart failure with preserved ejection fraction: a call for individualized therapies. *Eur Heart J* 2021;42:1940–1958.
16. Fraser AG, Tschöpe C, de Boer RA. Diagnostic recommendations and phenotyping for heart failure with preserved ejection fraction: knowing more and understanding less? *Eur J Heart Fail* 2021;23:964–972.
17. Raafs AG, Verdonschot JAJ, Henkens M, Adriaans BP, Wang P, Derks K, et al. The combination of carboxy-terminal propeptide of procollagen type I blood levels and late

- gadolinium enhancement at cardiac magnetic resonance provides additional prognostic information in idiopathic dilated cardiomyopathy – a multilevel assessment of myocardial fibrosis in dilated cardiomyopathy. *Eur J Heart Fail* 2021;23:933–944.
18. Sorimachi H, Obokata M, Takahashi N, Reddy YNV, Jain CC, Verbrugge FH, et al. Pathophysiologic importance of visceral adipose tissue in women with heart failure and preserved ejection fraction. *Eur Heart J* 2021;42:1595–1605.
 19. Withaar C, Meems LMG, de Boer RA. Fighting HFpEF in women: taking aim at belly fat. *Eur Heart J* 2021;42:1606–1608.
 20. Assmus B, Cremer S, Kirschbaum K, Culmann D, Kiefer K, Dorsheimer L, et al. Clonal haematopoiesis in chronic ischaemic heart failure: prognostic role of clone size for *DNMT3A*- and *TET2*-driver gene mutations. *Eur Heart J* 2021;42:257–265.
 21. Garnier S, Harakalova M, Weiss S, Mokry M, Regitz-Zagrosek V, Hengstenberg C, et al. Genome-wide association analysis in dilated cardiomyopathy reveals two new players in systolic heart failure on chromosomes 3p25.1 and 22q11.23. *Eur Heart J* 2021;42:2000–2011.
 22. Seferovic PM, Tsutsui H, McNamara DM, Ristic AD, Basso C, Bozkurt B, et al. Heart Failure Association of the ESC, Heart Failure Society of America and Japanese Heart Failure Society Position statement on endomyocardial biopsy. *Eur J Heart Fail* 2021;23:854–871.
 23. Tersalvi G, Dauw J, Gasperetti A, Winterton D, Cioffi GM, Scopigni F, et al. The value of urinary sodium assessment in acute heart failure. *Eur Heart J Acute Cardiovasc Care* 2021;10:216–223.
 24. Biegus J, Zymlinski R, Fudim M, Testani J, Sokolski M, Marciniak D, et al. Spot urine sodium in acute heart failure: differences in prognostic value on admission and discharge. *ESC Heart Fail* 2021;8:2597–2602.
 25. Mathew R, Di Santo P, Jung RG, Marbach JA, Hutson J, Simard T, et al. Milrinone as compared with dobutamine in the treatment of cardiogenic shock. *N Engl J Med* 2021;385:516–525.
 26. Karami M, Eriksen E, Ouweneel DM, Claessen BE, Vis MM, Baan J, et al. Long-term 5-year outcome of the randomized IMPRESS in severe shock trial: percutaneous mechanical circulatory support vs. intra-aortic balloon pump in cardiogenic shock after acute myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2021;10:1009–1015.
 27. Ceglarek U, Schellong P, Rosolowski M, Scholz M, Willenberg A, Kratzsch J, et al. The novel cystatin C, lactate, interleukin-6, and N-terminal pro-B-type natriuretic peptide (CLIP)-based mortality risk score in cardiogenic shock after acute myocardial infarction. *Eur Heart J* 2021;42:2344–2352.
 28. Arrigo M, Price S, Baran DA, Poss J, Aissaoui N, Bayes-Genis A, et al. Optimising clinical trials in acute myocardial infarction complicated by cardiogenic shock: a statement from the 2020 Critical Care Clinical Trialists Workshop. *Lancet Respir Med* 2021;9:1192–1202.
 29. Schmitto JD, Mariani S, Li T, Dogan G, Hanke JS, Bara C, et al. Five-year outcomes of patients supported with HeartMate 3: a single-centre experience. *Eur J Cardiothorac Surg* 2021;59:1155–1163.
 30. Cho S-M, Mehaffey JH, Myers SL, Cantor RS, Starling RC, Kirklin JK, et al. Cerebrovascular events in patients with centrifugal-flow left ventricular assist devices: a propensity score matched analysis from the intermacs registry. *Circulation* 2021;144:763–772.
 31. Kassner A, Oezpeker C, Gummert J, Zittermann A, Gartner A, Tiesmeier J, et al. Mechanical circulatory support does not reduce advanced myocardial fibrosis in patients with end-stage heart failure. *Eur J Heart Fail* 2021;23:324–334.
 32. Rivas-Lasarte M, Kumar S, Derbala MH, Ferrall J, Cefalu M, Rashid SMI, et al. Prediction of right heart failure after left ventricular assist implantation: external validation of the EUROMACS right-sided heart failure risk score. *Eur Heart J Acute Cardiovasc Care* 2021;10:723–732.
 33. Emerson D, Chikwe J, Catarino P, Hassanein M, Deng L, Cantor RS, et al. Contemporary left ventricular assist device outcomes in an aging population: an STS INTERMACS analysis. *J Am Coll Cardiol* 2021;78:883–894.
 34. Agbor-Enoh S, Shah P, Tunc I, Hsu S, Russell S, Feller E, et al. Cell-free DNA to detect heart allograft acute rejection. *Circulation* 2021;143:1184–1197.
 35. Sliwa K, van der Meer P, Petrie MC, Frogoudaki A, Johnson MR, Hilfiker-Kleiner D, et al. Risk stratification and management of women with cardiomyopathy/heart failure planning pregnancy or presenting during/after pregnancy: a position statement from the Heart Failure Association of the European Society of Cardiology Study Group on Peripartum Cardiomyopathy. *Eur J Heart Fail* 2021;23:527–540.
 36. Sliwa K, Petrie MC, van der Meer P, Mebazaa A, Hilfiker-Kleiner D, Jackson AM, et al. Clinical presentation, management, and 6-month outcomes in women with peripartum cardiomyopathy: an ESC EORP registry. *Eur Heart J* 2020;41:3787–3797.
 37. Farhan HA, Yaseen IF. Peripartum cardiomyopathy in Iraq: initial registry-based data and 6 month outcomes. *ESC Heart Fail* 2021;8:4048–4054.
 38. Mbakwem AC, Bauersachs J, Viljoen C, Hoevelmann J, van der Meer P, Petrie MC, et al. Electrocardiographic features and their echocardiographic correlates in peripartum cardiomyopathy: results from the ESC EORP PPCM registry. *ESC Heart Fail* 2021;8:879–889.
 39. Jackson AM, Petrie MC, Frogoudaki A, Laroche C, Gustafsson F, Ibrahim B, et al. Hypertensive disorders in women with peripartum cardiomyopathy: insights from the ESC EORP PPCM Registry. *Eur J Heart Fail* 2021. <https://doi.org/10.1002/ehf.2264>
 40. Spertus JA, Fine JT, Elliott P, Ho CY, Olivetto I, Saberi S, et al. Mavacamten for treatment of symptomatic obstructive hypertrophic cardiomyopathy (EXPLORER-HCM): health status analysis of a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet* 2021;397:2467–2675.
 41. Pelliccia F, Gersh BJ, Camici PG. Gaps in evidence for risk stratification for sudden cardiac death in hypertrophic cardiomyopathy. *Circulation* 2021;143:101–103.
 42. Marston NA, Han L, Olivetto I, Day SM, Ashley EA, Michels M, et al. Clinical characteristics and outcomes in childhood-onset hypertrophic cardiomyopathy. *Eur Heart J* 2021;42:1988–1996.
 43. Yilmaz A, Bauersachs J, Bengel F, Büchel R, Kindermann I, Klingel K, et al. Diagnosis and treatment of cardiac amyloidosis: position statement of the German Cardiac Society (DGK). *Clin Res Cardiol* 2021;110:479–506.
 44. Herrmann J, Lenihan D, Armenian S, Barac A, Blaes A, Cardinale D, et al. Defining cardiovascular toxicities of cancer therapies: an International Cardio-Oncology Society (IC-OS) consensus statement. *Eur Heart J* 2021. <https://doi.org/10.1093/eurheartj/ehab674>
 45. D'Souza M, Nielsen D, Svane IM, Iversen K, Rasmussen PV, Madelaire C, et al. The risk of cardiac events in patients receiving immune checkpoint inhibitors: a nationwide Danish study. *Eur Heart J* 2021;42:1621–1631.
 46. Totzeck M, Lutgens E, Neilan TG. Are we underestimating the potential for cardiotoxicity related to immune checkpoint inhibitors? *Eur Heart J* 2021;42:1632–1635.
 47. de Wit S, de Boer RA. From studying heart disease and cancer simultaneously to reverse cardio-oncology. *Circulation* 2021;144:93–95.
 48. Michel L, Helfrich I, Hendgen-Cotta UB, Mincu R-I, Korste S, Mroczek SM, et al. Targeting early stages of cardiotoxicity from anti-PD1 immune checkpoint inhibitor therapy. *Eur Heart J* 2021;ehab430.
 49. Lehmann LH, Cautela J, Palaskas N, Baik AH, Meijers WC, Allenbach Y, et al. Clinical strategy for the diagnosis and treatment of immune checkpoint inhibitor-associated myocarditis: a narrative review. *JAMA Cardiol* 2021;6:1329–1337.
 50. de Boer RA, Hulot J-S, Tocchetti CG, Aboumsallem JP, Ameri P, Anker SD, et al. Common mechanistic pathways in cancer and heart failure. A scientific roadmap on behalf of the Translational Research Committee of the Heart Failure Association (HFA) of the European Society of Cardiology (ESC). *Eur J Heart Fail* 2020;22:2272–2289.

51. Zannad F, Cotter G, Alonso Garcia A, George S, Davison B, Figtree G, et al. What can heart failure trialists learn from oncology trialists? *Eur Heart J* 2021;42:2373–2383.
52. Bauersachs J. Heart failure drug treatment: the fantastic four. *Eur Heart J* 2021;42:681–683.
53. McMurray JJV, Packer M. How should we sequence the treatments for heart failure and a reduced ejection fraction?: a redefinition of evidence-based medicine. *Circulation* 2021;143:875–877.
54. Rosano GMC, Moura B, Metra M, Bohm M, Bauersachs J, Ben Gal T, et al. Patient profiling in heart failure for tailoring medical therapy. A consensus document of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 2021;23:872–881.
55. Porcher R, Desguerre I, Amthor H, Chabrol B, Audic F, Rivier F, et al. Association between prophylactic angiotensin-converting enzyme inhibitors and overall survival in Duchenne muscular dystrophy—analysis of registry data. *Eur Heart J* 2021;42:1976–1984.
56. Bhatt AS, Vaduganathan M, Claggett BL, Liu J, Packer M, Desai AS, et al. Effect of sacubitril/valsartan vs. enalapril on changes in heart failure therapies over time: the PARADIGM-HF trial. *Eur J Heart Fail* 2021;23:1518–1524.
57. Tsutsui H, Momomura SI, Saito Y, Ito H, Yamamoto K, Sakata Y, et al. Efficacy and safety of sacubitril/valsartan in Japanese patients with chronic heart failure and reduced ejection fraction—results from the PARALLEL-HF study. *Circ J* 2021;85:584–594.
58. Proudfoot C, Studer R, Rajput T, Jindal R, Agrawal R, Corda S, et al. Real-world effectiveness and safety of sacubitril/valsartan in heart failure: a systematic review. *Int J Cardiol* 2021;331:164–171.
59. Giovannazzo S, Carmisciano L, Toma M, Benenati S, Tomasoni D, Sormani MP, et al. Sacubitril/valsartan in real-life European patients with heart failure and reduced ejection fraction: a systematic review and meta-analysis. *ESC Heart Fail* 2021;8:3547–3556.
60. Volpe M, Bauersachs J, Bayes-Genis A, Butler J, Cohen-Solal A, Gallo G, et al. Sacubitril/valsartan for the management of heart failure: a perspective viewpoint on current evidence. *Int J Cardiol* 2021;327:138–145.
61. Jackson AM, Jhund PS, Anand IS, Düngen HD, Lam CSP, Lefkowitz MP, et al. Sacubitril-valsartan as a treatment for apparent resistant hypertension in patients with heart failure and preserved ejection fraction. *Eur Heart J* 2021;42:3741–3752.
62. Felker GM, Butler J, Ibrahim NE, Piña IL, Maisel A, Bapat D, et al. Implantable cardioverter-defibrillator eligibility after initiation of sacubitril/valsartan in chronic heart failure: insights from PROVE-HF. *Circulation* 2021;144:180–182.
63. Docherty KF, Campbell RT, Brooksbank KJM, Dreisbach JG, Forsyth P, Godeseth RL, et al. Effect of neprilysin inhibition on left ventricular remodeling in patients with asymptomatic left ventricular systolic dysfunction late after myocardial infarction. *Circulation* 2021;144:199–209.
64. Docherty KF, Campbell RT, Brooksbank KJM, Godeseth RL, Forsyth P, McConnachie A, et al. Rationale and methods of a randomized trial evaluating the effect of neprilysin inhibition on left ventricular remodelling. *ESC Heart Fail* 2021;8:129–138.
65. Jering KS, Claggett B, Pfeiffer MA, Granger C, Køber L, Lewis EF, et al. Prospective ARNI vs. ACE inhibitor trial to Determine Superiority in reducing heart failure Events after Myocardial Infarction (PARADISE-MI): design and baseline characteristics. *Eur J Heart Fail* 2021;23:1040–1048.
66. Shah SJ, Cowie MR, Wachter R, Szecsy P, Shi V, Ibram G, et al. Baseline characteristics of patients in the PARALLAX trial: insights into quality of life and exercise capacity in heart failure with preserved ejection fraction. *Eur J Heart Fail* 2021;23:1541–1551.
67. Spertus JA. Quality of life in EMPEROR-Reduced: emphasizing what is important to patients while identifying strategies to support more patient-centred care. *Eur Heart J* 2021;42:1213–1215.
68. Kosiborod MN, Jhund PS, Docherty KF, Diez M, Petrie MC, Verma S, et al. Effects of dapagliflozin on symptoms, function, and quality of life in patients with heart failure and reduced ejection fraction: results from the DAPA-HF trial. *Circulation* 2020;141:90–99.
69. Butler J, Anker SD, Filippatos G, Khan MS, Ferreira JP, Pocock SJ, et al. Empagliflozin and health-related quality of life outcomes in patients with heart failure with reduced ejection fraction: the EMPEROR-Reduced trial. *Eur Heart J* 2021;42:1203–1212.
70. Abraham WT, Lindenfeld J, Ponikowski P, Agostoni P, Butler J, Desai AS, et al. Effect of empagliflozin on exercise ability and symptoms in heart failure patients with reduced and preserved ejection fraction, with and without type 2 diabetes. *Eur Heart J* 2021;42:700–710.
71. Solomon SD, Jhund PS, Claggett BL, Dewan P, Køber L, Kosiborod MN, et al. Effect of dapagliflozin in patients with HFrEF treated with sacubitril/valsartan: the DAPA-HF trial. *JACC Heart Fail* 2020;8:811–818.
72. Packer M, Anker SD, Butler J, Filippatos G, Ferreira JP, Pocock SJ, et al. Influence of neprilysin inhibition on the efficacy and safety of empagliflozin in patients with chronic heart failure and a reduced ejection fraction: the EMPEROR-Reduced trial. *Eur Heart J* 2021;42:671–680.
73. Lam CSP, Ferreira JP, Pfarr E, Sim D, Tsutsui H, Anker SD, et al. Regional and ethnic influences on the response to empagliflozin in patients with heart failure and a reduced ejection fraction: the EMPEROR-Reduced trial. *Eur Heart J* 2021;42:4442–4451.
74. Inzucchi SE, Docherty KF, Køber L, Kosiborod MN, Martinez FA, Ponikowski P, et al. Dapagliflozin and the incidence of type 2 diabetes in patients with heart failure and reduced ejection fraction: an exploratory analysis from DAPA-HF. *Diabetes Care* 2021;44:586–594.
75. McMurray JJV, Solomon SD, Docherty KF, Jhund PS. The Dapagliflozin and Prevention of Adverse outcomes in Heart Failure trial (DAPA-HF) in context. *Eur Heart J* 2021;42:1199–1202.
76. Anker SD, Butler J, Filippatos G, Ferreira JP, Bocchi E, Böhm M, et al. Empagliflozin in heart failure with a preserved ejection fraction. *N Engl J Med* 2021;385:1451–1461.
77. Anker SD, Butler J, Filippatos GS, Jamal W, Salsali A, Schnee J, et al. Evaluation of the effects of sodium–glucose co-transporter 2 inhibition with empagliflozin on morbidity and mortality in patients with chronic heart failure and a preserved ejection fraction: rationale for and design of the EMPEROR-Preserved Trial. *Eur J Heart Fail* 2019;21:1279–1287.
78. Solomon SD, de Boer RA, DeMets D, Hernandez AF, Inzucchi SE, Kosiborod MN, et al. Dapagliflozin in heart failure with preserved and mildly reduced ejection fraction: rationale and design of the DELIVER trial. *Eur J Heart Fail* 2021;23:1217–1225.
79. Bhatt DL, Szarek M, Steg PG, Cannon CP, Leiter LA, McGuire DK, et al. Sotagliflozin in patients with diabetes and recent worsening heart failure. *N Engl J Med* 2021;384:117–128.
80. Tromp J, Ponikowski P, Salsali A, Angermann CE, Biegus J, Blatchford J, et al. Sodium–glucose co-transporter 2 inhibition in patients hospitalized for acute decompensated heart failure: rationale for and design of the EMPULSE trial. *Eur J Heart Fail* 2021;23:826–834.
81. Heerspink HJL, Sjöström CD, Jongs N, Chertow GM, Kosiborod M, Hou FF, et al. Effects of dapagliflozin on mortality in patients with chronic kidney disease: a pre-specified analysis from the DAPA-CKD randomized controlled trial. *Eur Heart J* 2021;42:1216–1227.
82. Bhatt DL, Szarek M, Pitt B, Cannon CP, Leiter LA, McGuire DK, et al. Sotagliflozin in patients with diabetes and chronic kidney disease. *N Engl J Med* 2021;384:129–139.
83. Agarwal R, Kolkhof P, Bakris G, Bauersachs J, Haller H, Wada T, et al. Steroidal and non-steroidal mineralocorticoid receptor antagonists in cardiorenal medicine. *Eur Heart J* 2021;42:152–161.
84. Pitt B, Filippatos G, Agarwal R, Anker SD, Bakris GL, Rossing P, et al. Cardiovascular events with finerenone in kidney disease and type 2 diabetes. *N Engl J Med* 2021;385:2252–2263.
85. Cleland JGF, Ferreira JP, Mariotti B, Pellicori P, Cuthbert J, Verdonschot JAJ, et al. The effect of spironolactone on cardiovascular function and markers of fibrosis in people at increased risk of developing heart failure: the heart ‘OMICS’

- in AGEing (HOMAGE) randomized clinical trial. *Eur Heart J* 2021;42:684–696.
86. Ponikowski P, Alemyeh W, Oto A, Bahit MC, Noori E, Patel MJ, et al. Vericiguat in patients with atrial fibrillation and heart failure with reduced ejection fraction: insights from the VICTORIA trial. *Eur J Heart Fail* 2021;23:1300–1312.
 87. Voors AA, Mulder H, Reyes E, Cowie MR, Lassus J, Hernandez AF, et al. Renal function and the effects of vericiguat in patients with worsening heart failure with reduced ejection fraction: insights from the VICTORIA (Vericiguat Global Study in Subjects with HFrEF) trial. *Eur J Heart Fail* 2021;23:1313–1321.
 88. Teerlink JR, Diaz R, Felker GM, McMurray JJV, Metra M, Solomon SD, et al. Effect of ejection fraction on clinical outcomes in patients treated with omecamtiv mecarbil in GALACTIC-HF. *J Am Coll Cardiol* 2021;78:97–108.
 89. Jankowska EA, Kirwan BA, Kosiborod M, Butler J, Anker SD, McDonagh T, et al. The effect of intravenous ferric carboxymaltose on health-related quality of life in iron-deficient patients with acute heart failure: the results of the AFFIRM-AHF study. *Eur Heart J* 2021;42:3011–3020.
 90. Martens P, Dupont M, Dauw J, Nijst P, Herbots L, Dendale P, et al. The effect of intravenous ferric carboxymaltose on cardiac reverse remodelling following cardiac resynchronization therapy-the IRON-CRT trial. *Eur Heart J* 2021;ehab411.
 91. Tkaczyszyn M, Comin-Colet J, Voors AA, van Veldhuisen DJ, Enjuanes C, Moliner P, et al. Iron deficiency contributes to resistance to endogenous erythropoietin in anaemic heart failure patients. *Eur J Heart Fail* 2021;23:1677–1686.
 92. Täubel J, Hauke W, Rump S, Viereck J, Batkai S, Poetzsch J, et al. Novel antisense therapy targeting microRNA-132 in patients with heart failure: results of a first-in-human Phase 1b randomized, double-blind, placebo-controlled study. *Eur Heart J* 2021;42:178–188.
 93. Devaux Y, Badimon L. CDR132L: another brick in the wall towards the use of miRNAs to treat cardiovascular disease. *Eur Heart J* 2021;42:202–204.
 94. Baker AH, Giacca M. Antagonism of miRNA in heart failure: first evidence in human. *Eur Heart J* 2021;42:189–191.
 95. Hazebroek MR, Henkens MTHM, Raafs AG, Verdonchot JAJ, Merken JJ, Dennert RM, et al. Intravenous immunoglobulin therapy in adult patients with idiopathic chronic cardiomyopathy and cardiac parvovirus B19 persistence: a prospective, double-blind, randomized, placebo-controlled clinical trial. *Eur J Heart Fail* 2021;23:302–309.
 96. Brignole M, Pentimalli F, Palmisano P, Landolina M, Quartieri F, Occhetta E, et al. AV junction ablation and cardiac resynchronization for patients with permanent atrial fibrillation and narrow QRS: the APAF-CRT mortality trial. *Eur Heart J* 2021;42:4731–4739.
 97. Glikson M, Nielsen JC, Kronborg MB, Michowitz Y, Auricchio A, Barbash IM, et al. 2021 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy. *Europace*. 2021;euab232.
 98. Mullens W, Auricchio A, Martens P, Witte K, Cowie MR, Delgado V, et al. Optimized implementation of cardiac resynchronization therapy: a call for action for referral and optimization of care: A joint position statement from the Heart Failure Association (HFA), European Heart Rhythm Association (EHRA), and European Association of Cardiovascular Imaging (EACVI) of the European Society of Cardiology. *Eur J Heart Fail* 2020;22:2349–2369.
 99. Schrage B, Lund LH, Melin M, Benson L, Uijl A, Dahlstrom U, et al. Cardiac resynchronization therapy with or without defibrillator in patients with heart failure. *Europace* 2021;euab233.
 100. Otto CM, Nishimura RA, Bonow RO, Carabello BA, Erwin JP 3rd, Gentile F, et al. 2020 ACC/AHA guideline for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Thorac Cardiovasc Surg* 2021;162:e183–e353.
 101. Vahanian A, Beyersdorf F, Praz F, Milojevic M, Baldus S, Bauersachs J, et al. 2021 ESC/EACTS Guidelines for the management of valvular heart disease. *Eur Heart J* 2021;ehab395.
 102. Coats AJS, Anker SD, Baumbach A, Alfieri O, von Bardeleben RS, Bauersachs J, et al. The management of secondary mitral regurgitation in patients with heart failure: a joint position statement from the Heart Failure Association (HFA), European Association of Cardiovascular Imaging (EACVI), European Heart Rhythm Association (EHRA), and European Association of Percutaneous Cardiovascular Interventions (EAPCI) of the ESC. *Eur Heart J* 2021;42:1254–1269.
 103. Mack MJ, Lindenfeld J, Abraham WT, Kar S, Lim DS, Mishell JM, et al. 3-year outcomes of transcatheter mitral valve repair in patients with heart failure. *J Am Coll Cardiol* 2021;77:1029–1040.
 104. Kar S, Mack MJ, Lindenfeld J, Abraham WT, Asch FM, Weissman NJ, et al. Relationship between residual mitral regurgitation and clinical and quality-of-life outcomes after transcatheter and medical treatments in heart failure: COAPT trial. *Circulation* 2021;144:426–437.
 105. Gertz ZM, Herrmann HC, Lim DS, Kar S, Kapadia SR, Reed GW, et al. Implications of atrial fibrillation on the mechanisms of mitral regurgitation and response to MitraClip in the COAPT trial. *Circ Cardiovasc Interv* 2021;14:e010300.
 106. Jung B, Messika-Zeitoun D, Boutitie F, Trochu J-N, Armoiry X, Maucourt-Boulch D, et al. Characteristics and outcome of COAPT-eligible patients in the MITRA-FR trial. *Circulation* 2020;142:2482–2484.
 107. Lindenfeld J, Abraham WT, Grayburn PA, Kar S, Asch FM, Lim DS, et al. Association of effective regurgitation orifice area to left ventricular end-diastolic volume ratio with transcatheter mitral valve repair outcomes: a secondary analysis of the COAPT trial. *JAMA Cardiol* 2021;6:427–436.
 108. Lindenfeld J, Zile MR, Desai AS, Bhatt K, Ducharme A, Horstmannshof D, et al. Haemodynamic-guided management of heart failure (GUIDE-HF): a randomised controlled trial. *Lancet* 2021;398:991–1001.
 109. Bekfani T, Fudim M, Cleland JGF, Jorbenadze A, von Haehling S, Lorber A, et al. A current and future outlook on upcoming technologies in remote monitoring of patients with heart failure. *Eur J Heart Fail* 2021;23:175–185.
 110. Bozkurt B, Fonarow GC, Goldberg LR, Guglin M, Josephson RA, Forman DE, et al. Cardiac rehabilitation for patients with heart failure: JACC expert panel. *J Am Coll Cardiol* 2021;77:1454–1469.
 111. Kitzman DW, Whellan DJ, Duncan P, Pastva AM, Mentz RJ, Reeves GR, et al. Physical rehabilitation for older patients hospitalized for heart failure. *N Engl J Med* 2021;385:203–216.
 112. Harrison SL, Buckley BJR, Rivera-Caravaca JM, Zhang J, Lip GYH. Cardiovascular risk factors, cardiovascular disease, and COVID-19: an umbrella review of systematic reviews. *Eur Heart J Qual Care Clin Outcomes* 2021;7:330–339.
 113. Yoo J, Grewal P, Hotelling J, Papamanoli A, Cao K, Dhaliwal S, et al. Admission NT-proBNP and outcomes in patients without history of heart failure hospitalized with COVID-19. *ESC Heart Fail* 2021;8:4278–4287.
 114. Garg A, Seeliger B, Derda AA, Xiao K, Gietz A, Scherf K, et al. Circulating cardiovascular microRNAs in critically ill COVID-19 patients. *Eur J Heart Fail* 2021;23:468–475.
 115. Charman SJ, Velicki L, Okwose NC, Harwood A, McGregor G, Ristic A, et al. Insights into heart failure hospitalizations, management, and services during and beyond COVID-19. *ESC Heart Fail* 2021;8:175–182.
 116. Butt JH, Fosbøl EL, Gerds TA, Andersson C, Kragholm K, Biering-Sørensen T, et al. All-cause mortality and location of death in patients with established cardiovascular disease before, during, and after the COVID-19 lockdown: a Danish Nationwide Cohort Study. *Eur Heart J* 2021;42:1516–1523.
 117. Bauer A, Schreinlechner M, Sappner N, Dolejsi T, Tilg H, Aulinger BA, et al. Discontinuation versus continuation of renin-angiotensin-system inhibitors in COVID-19 (ACEI-COVID): a prospective, parallel group, randomised, controlled, open-label trial. *Lancet Respir Med* 2021;9:863–872.
 118. Lopes RD, Macedo AVS, de Barros ESPGM, Moll-Bernardes RJ, Dos Santos TM, Mazza L, et al. Effect of discontinuing vs continuing angiotensin-converting enzyme inhibitors and

- angiotensin II receptor blockers on days alive and out of the hospital in patients admitted with COVID-19: a randomized clinical trial. *JAMA* 2021;325:254–264.
119. Cohen JB, Hanff TC, William P, Sweitzer N, Rosado-Santander NR, Medina C, et al. Continuation versus discontinuation of renin–angiotensin system inhibitors in patients admitted to hospital with COVID-19: a prospective, randomised, open-label trial. *Lancet Respir Med* 2021;9:275–284.
 120. Kosiborod MN, Esterline R, Furtado RHM, Oscarsson J, Gasparyan SB, Koch GG, et al. Dapagliflozin in patients with cardiometabolic risk factors hospitalised with COVID-19 (DARE-19): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Diabetes Endocrinol* 2021;9:586–594.
 121. Bozkurt B, Kamat I, Hotez PJ. Myocarditis with COVID-19 mRNA vaccines. *Circulation* 2021;144:471–484.
 122. Huang C, Huang L, Wang Y, Li X, Ren L, Gu X, et al. 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study. *Lancet* 2021;397:220–232.
 123. Rajpal S, Tong MS, Borchers J, Zareba KM, Obarski TP, Simonetti OP, et al. Cardiovascular magnetic resonance findings in competitive athletes recovering from COVID-19 infection. *JAMA Cardiol* 2021;6:116–118.
 124. Clark DE, Parikh A, Dendy JM, Diamond AB, George-Durrett K, Fish FA, et al. COVID-19 myocardial pathology evaluation in athletes with cardiac magnetic resonance (COMPETE CMR). *Circulation* 2021;143:609–612.
 125. Martinez MW, Tucker AM, Bloom OJ, Green G, DiFiori JP, Solomon G, et al. Prevalence of inflammatory heart disease among professional athletes with prior COVID-19 infection who received systematic return-to-play cardiac screening. *JAMA Cardiol* 2021;6:745–752.
 126. Kawakami R, Sakamoto A, Kawai K, Gianatti A, Pellegrini D, Nasr A, et al. Pathological evidence for SARS-CoV-2 as a cause of myocarditis: JACC review topic of the week. *J Am Coll Cardiol* 2021;77:314–325.
 127. Kotecha T, Knight DS, Razvi Y, Kumar K, Vimalasvaran K, Thornton G, et al. Patterns of myocardial injury in recovered troponin-positive COVID-19 patients assessed by cardiovascular magnetic resonance. *Eur Heart J* 2021;42:1866–1878.